

## Tissue-Specific Regulation of the *WT1* Locus

Judith A. Hewitt, PhD, Patricia M. Kessler, MS, Christine E. Campbell, PhD, and  
Bryan R.G. Williams, PhD

The 11p13 Wilms' tumor locus consists of two coordinately regulated transcripts, *WT1* and *WIT-1*. These genes are highly expressed in the developing urogenital system, beginning with the urogenital ridge at day 10.5, the metanephric blastema at day 11.5, and during glomerular formation at day 13.5, becoming ultimately restricted to the podocytes. Stromal cells of the gonad also show abundant expression. *WT1* is expressed at lower levels in spleen, uterus, mesothelial linings of organs in the abdominal and thoracic cavities, and the ependymal layer of the ventral aspect of the spinal cord. *WIT-1* mRNA is about 10-fold less abundant than *WT1*, but appears to be expressed in the same tissue-restricted manner. Expression of the *WT1* protein is required for kidney development, although its physiological function remains to be determined. The function of *WIT-1* is similarly

unknown but one intriguing possibility is that it is an antisense regulator of *WT1*. An understanding of events controlling spatial and temporal regulation of these genes will greatly improve our ability to study the role of *WT1* and *WIT-1* in urogenital development. We have found that while chimeric reporter constructs containing 0.6–2.5 kb of the 5' region of the *WT1* gene direct transcription in many different cell lines, we were unable to detect expression in 13.5-day mouse embryos. However, a cosmid containing about 42 kb encompassing this region was able to direct the expression of abundant levels of mRNA from the appropriate transcription initiation sites in both stable transfectants of mouse Leydig cells (TM3) or in transgenic embryos. We are currently localizing the DNA elements required for this expression.

© 1996 Wiley-Liss, Inc.

**Key words:** Wilms' tumor, kidney development, transgenic mice

### INTRODUCTION

The *WT1* gene, originally identified on the basis of mutations and deletions in Wilms' tumors, has been shown to play a key role in urogenital development [1], consistent with its highly restricted pattern of spatial and temporal expression [2–6]. The *WT1* protein is a transcription factor, with four zinc fingers of the C<sub>2</sub>H<sub>2</sub> type [7]. Various splice forms and mutations have been identified, producing proteins with differing DNA binding activities in vitro [8–9]. From a developmental point of view, little is known about the function of *WT1*. Potential downstream targets of *WT1* have been identified by transfection studies in cell culture [10–14], but their role in kidney development has not been rigorously tested. Pax-2 has been identified as a target gene in vivo, as its expression is absent in metanephric blastema of homozygously deleted *WT1* <sup>-/-</sup> embryos [1,15].

Studies from our laboratory and others have shown that the region immediately surrounding the *WT1* transcriptional start site cluster, can confer transcriptional control on heterologous reporter genes in transfected cells in culture [16–20]. However, the promoter was expressed in many cell lines, including several which do not express the endogenous *WT1* gene [16,19]. An enhancer activity was identified in the 3' end of the *WT1* gene which is only active in K562 cells, and may play a role in the hematopoietic expression of *WT1* [19,21]. A silencer ac-

tivity was also detected within the third intron of the *WT1* gene which suppressed expression of the *WT1* promoter in nonrenal cell lines [22].

We have shown that the minimal promoter region, as defined in cell culture studies, will not direct appropriate expression of *WT1* in transgenic mouse embryos. However, we were able to demonstrate robust *WT1* expression from a 42-kilobase (kb) cosmid containing the first three exons of *WT1* plus ~30 kb of 5' sequence in both cultured cells and transgenic embryos. Efforts to localize enhancer elements within the 42-kb cosmid are currently in progress.

### MATERIALS AND METHODS

The murine *WT1* basal promoter was cloned by screening a genomic lambda fix II library prepared from 129/SV DNA (Stratagene, La Jolla, CA) with a 170-bp probe derived by polymerase chain reaction (PCR) from the 3' end of exon one of the murine Wilms' tumor (WT) gene representing nucleotides 750–920 [3]. Contiguous Hind

From the Department of Cancer Biology, Research Institute, The Cleveland Clinic Foundation, Cleveland, Ohio.

Received July 7, 1995; accepted February 6, 1996.

Address reprint requests to Bryan R.G. Williams, Ph.D., Department of Cancer Biology, Research Institute, Cleveland Clinic Foundation, 9500 Euclid Avenue, Cleveland OH 44195.

III and Eco RI fragments were isolated and cloned into the luciferase reporter vector pGL2-Basic (Promega, Madison, WI). Transient transfection assays were performed with Lipofectamine (GIBCO/BRL, Gaithersburg, MD). Expression of *WT1*/luciferase constructs in transient transfections of TM3 cells were assayed according to the manufacturer's instructions using the Reporter Lysis Buffer protocol (Promega).

The p $\beta$ GAL plasmid was constructed by blunting the 3.7-kb HindIII/BamHI fragment of pCH110 (Pharmacia) and inserting it into the EcoRV site of pBSII (Stratagene). The orientation was selected to be the same as the T7 promoter. The 2.5-kb PstI fragment of cosmid c2.2 (18) was cloned into the PstI site of p $\beta$ GAL, and the 1.3-kb SacI/PstI and 0.6 HindIII/PstI fragments were blunted and inserted into the Sma I site of p $\beta$ GAL. NotI/SalI digestion was used to purify the promoter/reporter fragment for transgenesis, and standard methods were used to produce FVB transgenic embryos [23]. Embryos were harvested at days 13.5–15.5 of development, fixed in 4% paraformaldehyde, cryosectioned (10  $\mu$ m) and stained for  $\beta$ -galactosidase activity overnight at 30°C and counterstained with neutral red [23]. Transgenic mice were identified by Southern blot analysis of placental/yolk sac DNA.

The cosmid c2.2 was transfected into TM3 cells (ATCC) by calcium phosphate precipitation or Lipofectamine (GIBCO/BRL). Stable transfectants were selected in 400 mg/ml G418 (GIBCO/BRL). RNA was prepared by the guanidinium isothiocyanate/CsCl gradient method and 20  $\mu$ g of total RNA was used for RNase protection using the 0.6 kb HindIII/PstI probe encompassing the *WT1* transcription start sites [18]. Transgenic embryos were prepared with cosmid c2.2 DNA linearized by the SalI site in the pWE15 vector. Total embryonic RNA was prepared by the RNazol B method (TelTest) and fifty  $\mu$ g was subjected to RNase protection as above.

BglII fragments of c2.2 were purified and inserted into the BamHI site of p1.3 $\beta$ GAL. Transfections and transgenic embryo analysis were performed as above.

## RESULTS

A comparison of the human and murine *WT1* loci shows extensive sequence homology 5' of the major *WT1* transcription initiation sites (Fig. 1A) [3]. Since minimal promoter activity for the human *WT1* gene has been established for fragments surrounding the transcription start site cluster, we determined whether this was also the case for the murine locus. The results (Fig. 1B) show that minimal bidirectional promoter activity can be assigned to a region in the murine locus homologous to the human counterpart.

To determine whether the minimal promoter defined in various cell lines would function in vivo in transgenic mouse embryos,  $\beta$ -galactosidase reporter constructs were

injected into one-cell mouse embryos, transferred to pseudopregnant recipients, and harvested between days 13.5 and 15.5 of gestation, counting the day of injection as day 0.5. This time point was chosen on the basis of maximal expression of endogenous *WT1* within the kidney. There was no  $\beta$ -galactosidase staining detected in any tissue examined in either transgenic embryos or their nontransgenic littermates in 0.6, 1.3 or 2.5  $\beta$ GAL embryos (data not shown). The results were obtained from 6 positive mice of 10 live births injected with the 2.5-kb promoter, 12 positive of 22 injected with the 1.3 kb promoter, and 3 positive of 7 injected with the 0.6 kb promoter. As similar constructs are reported to be active in various cell lines, yet are clearly not active in transgenic embryos, we concluded that in vivo expression of *WT1* requires additional regulatory sequences not required for transcription in cell cultures.


To identify those sequences, we performed transfection experiments with a cosmid, c2.2 (Fig. 2A), that contains the first three exons and introns of *WT1* as well as ~30 kb of 5' sequence that includes most if not all of the divergently transcribed *WIT-1* gene [24]. The major transcriptional start sites of *WT1* and *WIT-1* are about 600 bp apart, but additional *WIT-1* transcripts, which have not been completely mapped but which initiate within the *WT1* gene [18], are likely to be contained within this cosmid. The cosmid vector, pWE15, includes an SV40 driven *neo* gene, which was used in selecting stable transformants in TM3, a mouse Leydig cell line which expresses abundant endogenous *WT1* mRNA. RNase protection using a probe that spans the transcriptional start sites for *WT1* showed that 3 of 4 stable transfectants expressed significant amounts of human *WT1* mRNA. Moreover, this transcription was initiated at the same start sites utilized in human fetal kidney (Fig. 2B).

Cosmid c2.2 was also injected into embryos that were harvested on day 13.5 and used to prepare total embryonic RNA. RNase protection was performed with the same probe as in Figure 2A. Two of three transgenic embryos show abundant expression of *WT1* RNA from the transgene (Fig. 3), while there are no protected products in a nontransgenic littermate. Since total embryonic RNA was used in these experiments it is not possible to draw any conclusion about transcription start site usage, as it may be different in different tissues or aberrantly expressed in inappropriate tissues. However, the fact that we observe abundant expression of the transgene in both TM3 and transgenic embryos makes it possible to address these issues.


The observation that short promoter constructs (<2.5 kb) gave us very low reporter activity in cells, while longer constructs (42 kb) directed the expression of significant amounts of mRNA, prompted us to begin identification of regions in the cosmid that were responsible for this expression. BglII fragments of the cosmid, ranging

## A.


## Mouse



## Human



## B.




**Fig. 1. A:** Line drawing of mouse and human genomic fragments representing the Wilms' locus. The black boxes denote exons, the spaces between the boxes represent intron sequences. The transcriptional start sites are indicated by arrows. The arrow pointing to the left above the mouse sequence indicates antisense transcription of *WT1* as detected by RNase protection. The stippled bar indicates homology between mouse and human sequences over the marked region. A partial restriction map is represented as follows: E, EcoRI; P, PvuII; H, HindIII; B, BamHI. **B:** Transcriptional activity in transfected TM3 cells of a

segment of mouse DNA homologous to the human Wilms' putative bidirectional promoter. The constructs used are shown at the left with arrows indicating orientation of the 2.0-kb EcoRI fragment (see A). This fragment was cloned for use in the forward direction by blunt-end ligation into the SmaI site of pGL2-Basic (Promega, Madison, WI). The reverse orientation was cloned into the XhoI/BglII sites of the pGL2-Basic vector. The data are presented as luciferase activity after normalization for cotransfected  $\beta$ -galactosidase activity.

in size from 500 bp to 8 kb, were inserted upstream of the 1.3-kb minimal promoter in the  $\beta$ GAL constructs used above (Fig. 4). These were initially transfected into TM3 cells, as they had given robust activity in earlier experiments. Figure 4 shows that while most constructs had activity similar to the 1.3-kb minimal promoter, the addition of fragment I from BglII digestion of the cosmid enhanced transcriptional activity approximately fourfold.


Therefore, we concluded that the 2-kb fragment I had some transcriptional enhancing properties. This same construct was injected into mouse embryos as before and harvested on embryonic day 13.5. There were 10 live embryos, of which 3 were positive by Southern blotting of placental/yolk sac DNA. However, sectioning and staining revealed no  $\beta$ -galactosidase activity in the kidney or in other tissues (data not shown).



**Fig. 2.** Above: Map of cosmid and RNase protection probe. A schematic of the human DNA cosmid c2.2 is shown, with black boxes representing mapped exons. The major transcriptional start sites for *WT1* and *WIT-1* are shown, as well as the approximate location of a third transcript in this region, *WIT-3*. The line below the map indicates the position of the RNase protection probe used for *WT1* mRNA detection. Below: RNase protection of c2.2/TM3 stable transfectants. Twenty  $\mu$ g of total RNA was hybridized with the *WT1* probe, RNase digested and run on a denaturing gel. TM3-1 through TM3-4 are four different clonal lines. The sizes of protected products are indicated.

## DISCUSSION



Several reports have demonstrated transcriptional activity from 0.6 to 1.3 kb fragments of the promoter region of *WT1* in transfected cultured cells. We have referred to this activity as basal promoter activity [18], while others have referred to it as weak or promiscuous [16,19]. This level of activity was sufficient to demonstrate negative autoregulation by *WT1* overexpression in two studies [17,20]. Our conclusions from the experiments presented here are that this basal promoter region, while active in cell lines, is not active in transgenic embryos. We have chosen for analysis embryonic day 13.5, at which time endogenous *WT1* is expressed at high levels in a discrete pattern in the kidney and gonads, as well as at lower levels of expression in other tissues, such as spleen and mesothelium [6]. The results suggest that the identification of sequences required for the appropriate temporal



**Fig. 3.** Expression of c2.2 in transgenic embryos. Fifty  $\mu$ g of total embryonic RNA was hybridized with the same probe as in Figure 2. TM3-1 is the same clonal line as in Figure 2. Four embryos are presented here, indicated as positive or negative for the transgene. Hae III  $\phi$ x174 were kinased for markers.

and spatial regulation of the *WT1* locus in vivo will require the use of transgenic mouse embryos, in which all of the appropriate developmental signals are present.

Our efforts to identify these additional regulatory elements were initiated by using a 42-kb cosmid that contains the 5' end of the *WT1* gene (exons 1 to 3) and  $\sim$ 30-kb of 5' upstream sequence which includes most, if not all of the *WIT1* gene. Using this cosmid, we have demonstrated robust transgene expression both in the TM3 cell line and in transgenic mouse embryos. Moreover, as demonstrated by RNase protection, the correct transcriptional start sites are utilized in TM3 cells. To our knowledge this is the first demonstration of appropriate transcription start site usage in a transfected *WT1* promoter assay. It should be noted that this cosmid contains a *WT1* gene truncated within the third intron and therefore is missing



**Fig. 4.** Expression of  $\beta$ -galactosidase activity in TM3. Various BglIII fragments of c2.2, ranging in size from 8 to 0.5 kb, were subcloned into the p1.3 $\beta$ GAL vector as shown. These constructs were transfected transiently into TM3 cells using Lipofectamine and assayed for  $\beta$ -galactosidase activity. CMV $\beta$ GAL was used as a positive control for transfection. Dots represent duplicate samples.

its polyadenylation signal. We have not yet ascertained whether the transcripts being synthesized lack a polyA tail or are polyadenylated using a cryptic signal within the cosmid.

To identify the DNA elements in the cosmid responsible for transcriptional activity, we have made a series of reporter constructs consisting of the minimal 1.3-kb promoter fragment plus additional 0.5- to 8-kb fragments of the cosmid derived from sequences both 5' and 3' of the *WT1* transcriptional start sites. Having tested ~85% of the cosmid insert, no strong enhancer elements have been unequivocally identified. It is possible that the *WT1* enhancer(s) we seek lies in the remaining 15% of DNA still to be tested. Alternatively, the element may have been destroyed by the restriction digests used to dissect the cosmid. By digesting with a different restriction enzyme, we could perhaps keep intact such an enhancer. However, before pursuing these experiments, an alternative approach such as identifying DNase I hypersensitive sites within the transfected cosmid or endogenous murine *WT1* gene should provide important information in identifying the temporal and spatial elements regulating this locus.

## ACKNOWLEDGMENTS

This work was supported by USPHS Grant DK47101 from the National Institute of Diabetes and Digestive and Kidney Diseases.

## REFERENCES

- Kreidberg JA, Sariola H, Loring JM, Maeda M, Pelletier J, Housman D, Jaenisch R: *WT1* is required for early kidney development. *Cell* 74:679–691, 1993.
- Pritchard-Jones K, Fleming S, Davidson D, Bickmore W, Porteous D, Gosden C, Bard J, Buckler A, Pelletier J, Housman D, van Heyningen V, Hastie N: The candidate Wilms' tumour gene is involved in genitourinary development. *Nature* 346:194–197, 1990.
- Buckler AJ, Pelletier J, Haber DA, Glaser T, Housman DE: Isolation, characterization, and expression of the murine Wilms' tumor gene (*WT1*) during kidney development. *Mol Cell Biol* 11:1707–1712, 1991.
- Pelletier J, Schalling M, Buckler AJ, Rogers A, Haber DA, Housman D: Expression of the Wilms' tumor gene *WT1* in the murine urogenital system. *Genes Dev* 5:1345–1356, 1991.
- Armstrong JF, Pritchard-Jones K, Bickmore W, Hastie ND, Bard JBL: The expression of the Wilms' tumour gene, *WT1*, in the developing mammalian embryo. *Mech Dev* 40:85–97, 1992.
- Rackley RR, Flenniken AM, Kuriyan NP, Kessler PM, Stoler MH, Williams BRG: Expression of the Wilms' tumor suppressor gene *WT1* during mouse embryogenesis. *Cell Growth Diff* 4:1023–1031, 1993.
- Call KM, Glaser T, Ito CY, Buckler AJ, Pelletier J, Haber DA, Rose EA, Kral A, Yeger H, Lewis WH, Jones C, Housman DE: Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell* 60:509–520, 1990.
- Haber DA, Sohn RL, Buckler AJ, Pelletier J, Call KM, Housman DE: Alternative splicing and genomic structure of the Wilms tumor gene *WT1*. *Proc Natl Acad Sci* 88:9618–9622, 1991.
- Pelletier J, Bruening W, Kashtan CE, Mauer SM, Manivel JC, Striegel JE, Houghton DC, Junien C, Habib R, Fouser L, Fine RN, Silverman BL, Haber DA, Housman D: Germline mutations in the Wilms' tumor suppressor gene are associated with abnormal urogenital development in Denys-Drash syndrome. *Cell* 67:437–447, 1991.
- Drummond IA, Rupprecht HD, Rohwer-Nutter P, Lopez-Guisa JM, Madden SL, Rauscher FJ, Sukhatme VP: DNA recognition by splicing variants of the Wilms' tumor suppressor, *WT1*. *Mol Cell Biol* 14:3800–3809, 1994.
- Harrington MA, Konicek B, Song A, Xia XL, Fredericks WJ, Rauscher FJ: Inhibition of colony-stimulating factor-1 promoter activity by the product of the Wilms' tumor locus. *J Biol Chem* 268:21271–21275, 1993.
- Werner H, Re GG, Drummond IA, Sukhatme VP, Rauscher FJ, Sens DA, Garvin AJ, LeRoith D, Roberts CT: Increased expression of the insulin-like growth factor I receptor gene, *IGF1R*, in Wilms' tumor is correlated with modulation of *IGF1R* promoter activity by the *WT1* Wilms' tumor gene product. *Proc Natl Acad Sci* 90:5828–5832, 1993.
- Drummond IA, Madden SL, Rohwer-Nutter P, Bell GI, Sukhatme VP, Rauscher FJ: Repression of the insulin-like growth factor II gene by the Wilms' tumor suppressor *WT1*. *Science* 257:674–678, 1992.
- Dey BR, Sukhatme VP, Roberts AB, Sporn MB, Rauscher FJ, Kim SJ: Repression of the transforming growth factor- $\beta$ 1 gene by the Wilms' tumor suppressor *WT1* gene product. *Mol Endocrin* 8:595–602, 1994.
- Ryan G, Steele-Perkins V, Morris JF, Rauscher FJ, Dressler GR: Repression of *Pax-2* by *WT1* during normal kidney development. *Development* 121:867–875, 1995.
- Hofmann W, Royer HD, Drechsler M, Schneider S, Royer-Pokora B: Characterization of the transcriptional regulatory region of the human *WT1* gene. *Oncogene* 8:3123–3132, 1993.
- Rupprecht HD, Drummond IA, Madden SL, Rauscher FJ, Suk-

- hatme VP: The Wilms' tumor suppressor gene *WT1* is negatively autoregulated. *J Biol Chem* 269:6198–6206, 1994.
18. Campbell CE, Haung A, Gurney AL, Kessler PM, Hewitt JA, Williams BRG: Antisense transcripts and protein binding motifs within the Wilms' tumor (*WT1*) locus. *Oncogene* 9:583–595, 1994.
  19. Frazier GC, Wu YJ, Hewitt SM, Maity T, Ton CCT, Huff V, Saunders GF: Transcriptional regulation of the human Wilms' tumor gene (*WT1*). *J Biol Chem* 269:8892–8900, 1994.
  20. Malik KTA, Poirier V, Ivins SM, Brown KW: Autoregulation of the human *WT1* gene promoter. *FEBS Lett* 349:75–78, 1994.
  21. Wu YJ, Frazier GC, Saunders GF: GATA-1 transactivates the *WT1* hematopoietic specific enhancer. *J Biol Chem* 270:5944–5949, 1995.
  22. Frazier GC, Hewitt SM, Wu YJ, Maity T, Saunders GF: Identification and characterization of the regulatory regions of the human *WT1* gene in kidney and hematopoietic cells. *Proc Am Ass Cancer Res* 34:539, 1993.
  23. Hogan B, Beddington R, Constantini F, Lacy E: "Manipulating the Mouse Embryo." Cold Spring Harbor, NY: Cold Spring Harbor Press, 1994.
  24. Huang A, Campbell CE, Bonetta L, McAndrews-Hill MS, Chilton-MacNeill S, Coppes MJ, Law DJ, Feinberg AP, Yeger H, Williams BRG: Tissue, developmental, and tumor-specific expression of divergent transcripts in Wilms' tumor. *Science* 250:991–994, 1990.

## COMMENTARY

Hewitt et al. remind us of the possibility that a second gene, *WIT-1*, localized at chromosome 11p13 and very close to *WT1*, may play a role in urogenital development and Wilms' tumorigenesis. Although the function of *WIT-1* remains unknown, these investigators demonstrate that *WIT-1* mRNA expression which is about 10-fold less abundant than *WT1* expression, appears to be expressed in the same tissues and during the same stages of organ development as is *WT1*. The authors speculate that *WIT-1* might function as an antisense regulator of *WT1*.